


Pdgfrb

Rat Anti-Mouse PDGFR-beta (Clone 4C54) mAb

Catalog No.	CMP115	Quantity:	100 µg
Alternate Names:	Platelet-derived growth factor receptor beta, PDGFR-beta, CD140 antigen-like family member B, CD140b, Platelet-derived growth factor receptor 1, PDGFR-1		
Description:	<p>PDGFRα and Rβ are members of the class III subfamily of receptor tyrosine kinases (RTK) that also includes the receptors for MCSF, SCF, and Flt3 ligand. All class III RTKs are characterized by the presence of five immunoglobulin-like domains in their extracellular region and a split kinase domain in their intracellular region. Ligand-induced receptor dimerization results in autophosphorylation in trans resulting in the activation of several intracellular signaling pathways that can lead to cell proliferation, cell survival, cytoskeletal rearrangement, and cell migration. Many cell types, including fibroblasts and smooth muscle cells, express both the α and β receptors. Others have only the α receptors (oligodendrocyte progenitor cells, mesothelial cells, liver sinusoidal endothelial cells, astrocytes, platelets, and megakaryocytes) or only the β receptors (myoblasts, capillary endothelial cells, pericytes, T cells, myeloid hematopoietic cells, and macrophages). Recombinant mouse and human soluble PDGF Rβ bind PDGF with high affinity and are potent PDGF antagonists.</p>		
UniProt ID:	P05622		
Gene ID:	18596		
Source:	Cell culture supernatant		
Specificity:	Recognizes mouse PDGFR-beta. No cross-reactivity with mouse PDGF receptor-alpha.		
Immunogen:	Recombinant mouse PDGFR-beta extracellular domain		
Isotype:	Rat IgG2		
Clone:	4C54		
Formulation:	Lyophilized from sterile filtered PBS		
Purification:	Protein G affinity chromatography		
Reconstitution:	<p>Centrifuge vial briefly prior to opening. Add 500 µl sterile PBS to the vial for a concentration of 200 µg/ml. Gently mix, allow a few minutes to solubilize completely. Do not vortex.</p>		
Cross-Reactivity:	Reactivity to other species has not been tested.		
Applications:	<p>Western Blot: suggested dilution 1:500-1:1000 IHC-P: suggested dilution 1:50-1:200</p>		
Storage & Stability:	<p>Lyophilized antibody is stable for at least 1 year from date of receipt at -20°C to -80°C. Reconstituted antibody may be prepared as aliquots and stored at -20°C to -80°C for at least six months. Avoid repeated freeze-thaw cycles.</p>		

PDGFR-beta IHC of paraffin sections of mouse pancreas from LPS exposed animals.



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com