


CCL3

Recombinant Human MIP-1 alpha / CCL3

Catalog No.	CRM400A CRM400B CRM400C	Quantity:	5 µg 100 µg 1.0 mg
Alternate Names:	Macrophage Inflammatory Protein 1-alpha, CCL3, LD78-alpha, MIP1A, MIP-1-alpha, SCI, SCYA3, SIS-beta, TY5		
Description:	Both MIP-1 alpha and MIP-1 beta are structurally and functionally related by having the four highly conserved cysteine residues present in CC chemokines. They participate in the host response to invading bacterial, viral, parasite and fungal pathogens by regulating the trafficking and activation state of selected subgroups of inflammatory cells e.g. macrophages, lymphocytes and NK cells. While both MIP-1 alpha and MIP-1 beta exert similar effects on monocytes their effect on lymphocytes differ; with MIP-1 alpha selectively attracting CD8+ lymphocytes and MIP-1 beta selectively attracting CD4+ lymphocytes. Additionally, MIP-1 alpha and MIP-1 beta have also been shown to be potent chemoattractants for B cells, eosinophils and dendritic cells. Both human and murine MIP-1 alpha and MIP-1 beta are active on human and murine hematopoietic cells.		
UniProt ID:	P10147		
Gene ID:	6348		
Source:	<i>E. coli</i>		
Molecular Weight:	~7.8 kDa (70 aa)		
Formulation:	Lyophilized from sterile-filtered 20 mM PB, pH 7.4, containing 100 mM NaCl.		
Purity:	>96% as determined by HPLC and SDS-PAGE analyses.		
Endotoxin Level:	<1 EU/µg of recombinant human MIP-1α/CCL3 as determined by LAL method.		
Biological Activity:	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human monocytes is in a concentration range of 1.0-10 ng/ml.		
Amino Acid Sequence:	ASLAADTPA CCFSYTSRQI PQNFIADYFE TSSQCCKPGV IFLTKRSRQV CADPSEEWVQ KYVSDLELSA		
Reconstitution:	Centrifuge vial prior to opening. Add sterile distilled water or aqueous buffer to a concentration of 0.1-1.0 mg/ml. Further dilutions should be made in appropriate buffered solutions.		
Storage & Stability:	Upon receipt, store at -20°C to -80°C for up to 1 year. After reconstitution, the preparation is stable for up to one week at 2-8°C or store in aliquots at -20°C to -80°C. For long term storage of reconstituted protein, it is recommended that a carrier protein such as 0.1% BSA or HSA be added. This depends on the particular application. Avoid repeated freeze/thaw cycles.		





NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com