


TF

Native Human Apotransferrin

Catalog No.	CSI19707A CSI19707B	Quantity:	100 mg 1 g
Alternate Names:	Serotransferrin, beta-1-metal-binding globulin, siderophilin, TRF, apo-hTF, apotransferrin		
Description:	<p>Transferrin (TF) is a glycoprotein thought to have been created as a result of an ancient gene duplication event that led to generation of homologous C- and N-terminal domains each of which binds one ion of Ferric Iron. Therefore, each Human TF molecule has the ability to carry two Iron ions in the Ferric form (Fe³⁺). The function of TF is to transport Iron from the intestine, reticuloendothelial system, and liver parenchymal cells to all proliferating cells in the body. Human HOLO transferrin is iron saturated whereas APO transferrin is substantially iron-free. When human transferrin loaded with iron encounters a human transferrin receptor on the surface of a cell, it binds to it and is consequently transported into the cell in a vesicle. The cell will acidify the vesicle, causing human transferrin to release its iron ions.</p> <p>Apotransferrin is an important transport factor used in defined culture media. Purified to have less than 0.01% iron and an iron-binding capacity of 1.25 ug/mg. Dissolves in water at 10 mg/ml.</p> <p>Low endotoxin, bioprocessing grade in bulk quantity is available.</p>		
UniProt ID	P02787		
Gene ID:	7018		
Source:	Human plasma		
Iron Content:	< 30 µg per gram		
Molecular Weight:	80 kDa		
Formulation:	Lyophilized from 10 mM Na phosphate buffer, pH 7.4		
Purity:	≥ 98% by SDS-PAGE		
Reconstitution:	Centrifuge vial prior to opening. Add sterile distilled water to a concentration of > 0.1 mg/ml and gently pipette the solution up and down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution.		
Storage & Stability:	Store lyophilized protein at -20 to -80°C for at least 1 year or reconstituted stock solution in aliquots at -20°C to -80°C. Avoid repeated freeze-thaw cycles.		
Infectious Disease Statement:	Prepared from plasma shown to be non reactive for HBsAg, anti-HCV, anti-HBc, and negative for anti-HIV 1 & 2 by FDA approved tests.		



SDS-PAGE: 4 - 12% Bis-Tris NuPAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com