

## Mouse Anti-Human CD15 (Clone MEM-158) mAb

<b>Catalog No.</b>	MON1182	<b>Quantity:</b>	100 µg
<b>Alternate Names:</b>	Lewis X, Le(x) stage-specific embryonic antigen-1, SSEA-1, 3- fucosyl-N-acetyl-lactosamine		
<b>Description:</b>	<p>CD15 (Lewis X, Le(x) stage-specific embryonic antigen-1, SSEA-1) is a trisaccharide determinant (3- fucosyl-N-acetyllactosamine) expressed on several glycolipids, glycoproteins and proteoglycans of various cell types, e.g. granulocytes, mast cells, monocytes, macrophages, cells of gastric mucosa, nervous system or various tumor cells, expressed on virtually all peripheral blood granulocytes. The CD15 antigen is expressed on approximately 95% of human peripheral blood eosinophils and neutrophils and is present on approximately 20% of mature circulating monocytes. Anti-CD15 has virtually no reactivity with T or B cells.</p> <p>There are several variants of Lewis x, such as sialyl-Lewis x or sulphated Lewis x. Cells with high surface expression of Le(x) antigen exhibit strong self-aggregation, based on calcium-dependent Le(x)-Le(x) interaction. This process is involved for example in embryo compaction or in autoaggregation of teratocarcinoma cells. Sialyl Le(x) and its isomer sialyl-Le(a) are ligands of selectins. CD15 expression has been extensively used to confirm diagnosis of Hodgkin and acute disease.</p>		
<b>Concentration:</b>	1 mg/ml		
<b>Specificity:</b>	Human CD15		
<b>Host:</b>	Mouse		
<b>Immunogen:</b>	Human granulocytes		
<b>Isotype:</b>	IgM		
<b>Clone:</b>	MEM-158		
<b>Formulation:</b>	Tris-buffered saline (TBS) containing 15 mM sodium azide		
<b>Applications:</b>	IP, FC		
<b>Storage &amp; Stability:</b>	Store undiluted antibody at 2-8°C until expiration date.		

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.