

Human Jurkat Cell Lysate

Catalog No. PX318A **Quantity:** 100 µg

Description: Jurkat cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl fluoride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% β-mercaptoethanol.

Cell Line: Jurkat, clone E6-1 (Human acute T cell leukemia)

Growth Media: RPMI with 10% NCS (Newborn calf serum)

Concentration: 2 mg/ml

Formulation: Lysate is supplied in 1 x SDS sample buffer containing 5% β-mercaptoethanol.

Applications: **Centrifuge vial before use.** Jurkat cell lysate is ready to load on SDS-PAGE for Western blotting, 10-20 µg per lane is recommended for mini gel.

Storage & Stability: Store at 2-8°C for continuous use. Store in working aliquots at -80°C for up to 1 year. **Avoid repeated freeze-thaw cycles.**

Certification: This material has been tested by accepted techniques and has been found to be negative for HBsAg, HIV 1/2, and HCV.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com