

## MDA-MB-361 Cell Lysate

**Catalog No.** PX322A **Quantity:** 100 µg

**Description:** MDA-MB-361 cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl fluoride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% β-mercaptoethanol.

**Concentration:** 2 mg/mL

**Cell Line:** MDA-MB-361 (Human Breast Adenocarcinoma)

**Growth Media:** DMEM/Hams F12 with 10% NCS (Newborn Calf serum)

**Formulation:** Lysate is supplied as whole cell lysate in 1 x SDS sample buffer containing 5% β-mercaptoethanol.

**Applications:** **Centrifuge vial before use.** MDA-MB-361 cell lysate is ready to load on SDS-PAGE for Western blotting, 10-20 µg per lane is recommended for mini gel.

**Storage & Stability:** Store at 2-8°C for continuous use. Store in working aliquots at -80°C for up to 1 year. **Avoid repeated freeze-thaw cycles.**

**Certification:** This material has been tested by accepted techniques and has been found to be negative for HBsAg, HIV 1/2, and HCV.

**NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.**



**Cell Sciences®**  
65 Parker Street  
Unit 11  
Newburyport, MA 01950

Toll Free: 888-769-1246  
Phone: 978-572-1070  
Fax: 978-992-0298

E-mail: [info@cellsciences.com](mailto:info@cellsciences.com)  
Website: [www.cellsciences.com](http://www.cellsciences.com)